

PERFORMANCE HI JACK R20

Medicated

Beef Cattle - Feedlot Supplement

For improved feed efficiency in cattle fed in confinement for slaughter
ACTIVE DRUG INGREDIENTS

MONENSIN 30 g/ton

GUARANTEED ANALYSIS

CRUDE PROTEIN, min.....12.0 %	SALT, min.....0.1 %
CRUDE FAT, min.....2.0 %	SALT, max.....0.6 %
CRUDE FIBER, max.....18.0 %	POTASSIUM, min.....0.7%
CALCIUM, min.....1.2 %	VITAMIN A, min.....2,000 IU/lb
CALCIUM, max.....1.7 %	VITAMIN D, min.....200 IU/lb
PHOSPHORUS, min.....0.4 %	VITAMIN E, min.....2.5 IU/lb

INGREDIENTS

Processed grain by-products, roughage products, grain products, calcium carbonate, molasses products, sodium bicarbonate, salt, pork fat (preserved with BHA), zinc sulfate, manganese sulfate, ferrous sulfate, vitamin A supplement, vitamin D supplement, copper sulfate, copper chloride, sodium selenite, ethylenediamine dihydroiodide, cobalt carbonate, vitamin E supplement, thiamine mononitrate

WARNING: This product, which contains added copper, should not be fed to sheep or related species that have a low tolerance to copper. This product has been formulated specifically for swine and is not intended for other species.

FEEDING DIRECTIONS:

Each pound of Performance Hi Jack R20 contains 10 milligrams of Monensin. Feed between 5 to 30 pounds of Performance Hi Jack R20 per head per day. Feed continuously with a forage source to provide each beef animal with not less than 50 nor more than 300 milligrams Monensin per head per day. Always provide clean, fresh water.

CAUTION: Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and the amount of refusals fed should be taken into consideration to prevent Monensin overdosing. **WARNING:** A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

This feed does not contain urea, ruminant meat and bone meal, chicken litter, or restaurant grease.

Net weight 50 lb. (22.68 kg) or bulk invoice

MANUFACTURED BY:

PERFORMANCE FEEDS

2784 Bethany Road
Red Boiling Springs, TN 37150
(866) 699-5080
(615) 699-5080

LoadSheet #



PERFORMANCE HI JACK R20

Medicated

Beef Cattle - Feedlot Supplement

For improved feed efficiency in cattle fed in confinement for slaughter
ACTIVE DRUG INGREDIENTS

MONENSIN 30 g/ton

GUARANTEED ANALYSIS

CRUDE PROTEIN, min.....12.0 %	SALT, min.....0.1 %
CRUDE FAT, min.....2.0 %	SALT, max.....0.6 %
CRUDE FIBER, max.....18.0 %	POTASSIUM, min.....0.7%
CALCIUM, min.....1.2 %	VITAMIN A, min.....2,000 IU/lb
CALCIUM, max.....1.7 %	VITAMIN D, min.....200 IU/lb
PHOSPHORUS, min.....0.4 %	VITAMIN E, min.....2.5 IU/lb

INGREDIENTS

Processed grain by-products, roughage products, grain products, calcium carbonate, molasses products, sodium bicarbonate, salt, pork fat (preserved with BHA), zinc sulfate, manganese sulfate, ferrous sulfate, vitamin A supplement, vitamin D supplement, copper sulfate, copper chloride, sodium selenite, ethylenediamine dihydroiodide, cobalt carbonate, vitamin E supplement, thiamine mononitrate

WARNING: This product, which contains added copper, should not be fed to sheep or related species that have a low tolerance to copper. This product has been formulated specifically for swine and is not intended for other species.

FEEDING DIRECTIONS:

Each pound of Performance Hi Jack R20 contains 10 milligrams of Monensin. Feed between 5 to 30 pounds of Performance Hi Jack R20 per head per day. Feed continuously with a forage source to provide each beef animal with not less than 50 nor more than 300 milligrams Monensin per head per day. Always provide clean, fresh water.

CAUTION: Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and the amount of refusals fed should be taken into consideration to prevent Monensin overdosing. **WARNING:** A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

This feed does not contain urea, ruminant meat and bone meal, chicken litter, or restaurant grease.

Net weight 50 lb. (22.68 kg) or bulk invoice

MANUFACTURED BY:

PERFORMANCE FEEDS

2784 Bethany Road
Red Boiling Springs, TN 37150
(866) 699-5080
(615) 699-5080

LoadSheet #



PERFORMANCE HI JACK R20

Medicated

Beef Cattle - Feedlot Supplement

For improved feed efficiency in cattle fed in confinement for slaughter
ACTIVE DRUG INGREDIENTS

MONENSIN 30 g/ton

GUARANTEED ANALYSIS

CRUDE PROTEIN, min.....12.0 %	SALT, min.....0.1 %
CRUDE FAT, min.....2.0 %	SALT, max.....0.6 %
CRUDE FIBER, max.....18.0 %	POTASSIUM, min.....0.7%
CALCIUM, min.....1.2 %	VITAMIN A, min.....2,000 IU/lb
CALCIUM, max.....1.7 %	VITAMIN D, min.....200 IU/lb
PHOSPHORUS, min.....0.4 %	VITAMIN E, min.....2.5 IU/lb

INGREDIENTS

Processed grain by-products, roughage products, grain products, calcium carbonate, molasses products, sodium bicarbonate, salt, pork fat (preserved with BHA), zinc sulfate, manganese sulfate, ferrous sulfate, vitamin A supplement, vitamin D supplement, copper sulfate, copper chloride, sodium selenite, ethylenediamine dihydroiodide, cobalt carbonate, vitamin E supplement, thiamine mononitrate

WARNING: This product, which contains added copper, should not be fed to sheep or related species that have a low tolerance to copper. This product has been formulated specifically for swine and is not intended for other species.

FEEDING DIRECTIONS:

Each pound of Performance Hi Jack R20 contains 10 milligrams of Monensin. Feed between 5 to 30 pounds of Performance Hi Jack R20 per head per day. Feed continuously with a forage source to provide each beef animal with not less than 50 nor more than 300 milligrams Monensin per head per day. Always provide clean, fresh water.

CAUTION: Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and the amount of refusals fed should be taken into consideration to prevent Monensin overdosing. **WARNING:** A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

This feed does not contain urea, ruminant meat and bone meal, chicken litter, or restaurant grease.

Net weight 50 lb. (22.68 kg) or bulk invoice

MANUFACTURED BY:

PERFORMANCE FEEDS
2784 Bethany Road
Red Boiling Springs, TN 37150
(866) 699-5080
(615) 699-5080

LoadSheet #

