

**PERFORMANCE GROWER  
SUPREME 14  
CATTLE RATION R20  
MEDICATED**

**BEEF CATTLE—GROWER**

For improved feed efficiency in cattle fed in confinement for slaughter.

**ACTIVE DRUG INGREDIENT**

MONENSIN . . . . . 20 g/ton

**GUARANTEED ANALYSIS**

CRUDE PROTEIN, min . . . . . 14.0%	PHOSPHORUS (P), min . . . . . 0.60%
* This includes not more than 1.0% equivalent protein from non-protein nitrogen.	SALT (NaCl), min . . . . . 0.2%
CRUDE FAT, min . . . . . 3.0%	SALT (NaCl), max . . . . . 0.7%
CRUDE FIBER, max . . . . . 18.0%	POTASSIUM (K), min . . . . . 0.6%
CALCIUM (Ca.), min . . . . . 1.3%	VITAMIN A, min . . . . . 3000 IU/lb
CALCIUM (Ca.), max . . . . . 1.8%	VITAMIN D3, min . . . . . 300 IU/lb

**INGREDIENTS**

Forage products, processed grain by-products, plant protein products, urea, salt, calcium carbonate, sodium selenite, zinc sulfate, manganese sulfate, copper sulfate, copper chloride, ferrous sulfate, ethylenediamine dihydrochloride, cobalt carbonate, vitamin A acetate, vitamin D supplement, thiamine mononitrate, vitamin E supplement.

**WARNING:** This product, which contains added copper, should not be fed to sheep or related species that have a low tolerance to copper.

**CAUTION: USE AS DIRECTED**

This product has been formulated specifically for beef and is not intended for other species.

**FEEDING DIRECTIONS:**

Each pound of Grower Supreme 14 Cattle Ration R20 contains 10 milligrams of Monensin. Feed between 5 to 30 pounds of Grower Supreme 14 Cattle Ration R20 per head per day. Feed continuously with a forage source to provide each beef animal with not less than 50 nor more than 360 milligrams Monensin per head per day. Always provide clean, fresh water.

**CAUTION:** Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and the amount of refusals fed should be taken into consideration to prevent Monensin overdosing. A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

This feed does not contain ruminant meat and bone meal, chicken litter, or restaurant grease. Seller warrants only that this feed conforms to the description on the label and is suitable for the purposes set forth when used in accordance with the directions for use on said label.

Net weight 50 lb. (22.68 kg) or bulk invoice

**MANUFACTURED BY:  
PERFORMANCE FEEDS**

2784 Bethany Road  
Red Boiling Springs, TN 37150  
(866) 699-5080  
(615) 699-5080



LoadSheet #

**PERFORMANCE GROWER  
SUPREME 14  
CATTLE RATION R20  
MEDICATED**

**BEEF CATTLE—GROWER**

For improved feed efficiency in cattle fed in confinement for slaughter.

**ACTIVE DRUG INGREDIENT**

MONENSIN . . . . . 20 g/ton

**GUARANTEED ANALYSIS**

CRUDE PROTEIN, min . . . . . 14.0%	PHOSPHORUS (P), min . . . . . 0.60%
* This includes not more than 1.0% equivalent protein from non-protein nitrogen.	SALT (NaCl), min . . . . . 0.2%
CRUDE FAT, min . . . . . 3.0%	SALT (NaCl), max . . . . . 0.7%
CRUDE FIBER, max . . . . . 18.0%	POTASSIUM (K), min . . . . . 0.6%
CALCIUM (Ca.), min . . . . . 1.3%	VITAMIN A, min . . . . . 3000 IU/lb
CALCIUM (Ca.), max . . . . . 1.8%	VITAMIN D3, min . . . . . 300 IU/lb

**INGREDIENTS**

Forage products, processed grain by-products, plant protein products, urea, salt, calcium carbonate, sodium selenite, zinc sulfate, manganese sulfate, copper sulfate, copper chloride, ferrous sulfate, ethylenediamine dihydrochloride, cobalt carbonate, vitamin A acetate, vitamin D supplement, thiamine mononitrate, vitamin E supplement.

**WARNING:** This product, which contains added copper, should not be fed to sheep or related species that have a low tolerance to copper.

**CAUTION: USE AS DIRECTED**

This product has been formulated specifically for beef and is not intended for other species.

**FEEDING DIRECTIONS:**

Each pound of Grower Supreme 14 Cattle Ration R20 contains 10 milligrams of Monensin. Feed between 5 to 30 pounds of Grower Supreme 14 Cattle Ration R20 per head per day. Feed continuously with a forage source to provide each beef animal with not less than 50 nor more than 360 milligrams Monensin per head per day. Always provide clean, fresh water.

**CAUTION:** Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and the amount of refusals fed should be taken into consideration to prevent Monensin overdosing. A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

This feed does not contain ruminant meat and bone meal, chicken litter, or restaurant grease. Seller warrants only that this feed conforms to the description on the label and is suitable for the purposes set forth when used in accordance with the directions for use on said label.

Net weight 50 lb. (22.68 kg) or bulk invoice

**MANUFACTURED BY:  
PERFORMANCE FEEDS**

2784 Bethany Road  
Red Boiling Springs, TN 37150  
(866) 699-5080  
(615) 699-5080



LoadSheet #

**PERFORMANCE GROWER  
SUPREME 14  
CATTLE RATION R20  
MEDICATED**

**BEEF CATTLE—GROWER**

For improved feed efficiency in cattle fed in confinement for slaughter.

**ACTIVE DRUG INGREDIENT**

MONENSIN . . . . . 20 g/ton

**GUARANTEED ANALYSIS**

CRUDE PROTEIN, min . . . . . 14.0%	PHOSPHORUS (P), min . . . . . 0.60%
* This includes not more than 1.0% equivalent protein from non-protein nitrogen.	SALT (NaCl), min . . . . . 0.2%
CRUDE FAT, min . . . . . 3.0%	SALT (NaCl), max . . . . . 0.7%
CRUDE FIBER, max . . . . . 18.0%	POTASSIUM (K), min . . . . . 0.6%
CALCIUM (Ca.), min . . . . . 1.3%	VITAMIN A, min . . . . . 3000 IU/lb
CALCIUM (Ca.), max . . . . . 1.8%	VITAMIN D3, min . . . . . 300 IU/lb

**INGREDIENTS**

Forage products, processed grain by-products, plant protein products, urea, salt, calcium carbonate, sodium selenite, zinc sulfate, manganese sulfate, copper sulfate, copper chloride, ferrous sulfate, ethylenediamine dihydrochloride, cobalt carbonate, vitamin A acetate, vitamin D supplement, thiamine mononitrate, vitamin E supplement.

**WARNING:** This product, which contains added copper, should not be fed to sheep or related species that have a low tolerance to copper.

**CAUTION: USE AS DIRECTED**

This product has been formulated specifically for beef and is not intended for other species.

**FEEDING DIRECTIONS:**

Each pound of Grower Supreme 14 Cattle Ration R20 contains 10 milligrams of Monensin. Feed between 5 to 30 pounds of Grower Supreme 14 Cattle Ration R20 per head per day. Feed continuously with a forage source to provide each beef animal with not less than 50 nor more than 360 milligrams Monensin per head per day. Always provide clean, fresh water.

**CAUTION:** Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and the amount of refusals fed should be taken into consideration to prevent Monensin overdosing. A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

This feed does not contain ruminant meat and bone meal, chicken litter, or restaurant grease. Seller warrants only that this feed conforms to the description on the label and is suitable for the purposes set forth when used in accordance with the directions for use on said label.

Net weight 50 lb. (22.68 kg) or bulk invoice

**MANUFACTURED BY:  
PERFORMANCE FEEDS**

2784 Bethany Road  
Red Boiling Springs, TN 37150  
(866) 699-5080  
(615) 699-5080



LoadSheet #