

PERFORMANCE ELITE CREEP FEED CATTLE RATION R40

MEDICATED

Calves-Birth to Weaning

Increased rate of weight gain in cattle (pastured slaughter, stocker, and feeder cattle, and dairy and beef replacement heifers).

ACTIVE DRUG INGREDIENT

MONENSIN.....40 g/ton

GUARANTEED ANALYSIS

CRUDE PROTEIN, min... 15.0 %	SALT, min.....0.1 %
CRUDE FAT, min..... 3.0 %	SALT, max..... 0.6 %
CRUDE FIBER, max..... 16.0 %	POTASSIUM, min.....0.7 %
CALCIUM, min.....1.1 %	VITAMIN A, min.....4,000 IU/lb
CALCIUM, max.....1.6 %	VITAMIN D, min.....400 IU/lb
PHOSPHORUS, min.....0.4 %	VITAMIN E, min.....5 IU/lb

INGREDIENTS

Processed grain by-products, roughage products, grain products, calcium carbonate, molasses products, salt, pork fat (preserved with BHA), zinc sulfate, manganese sulfate, ferrous sulfate, vitamin A supplement, vitamin D supplement, copper chloride, copper sulfate, sodium selenite, ethylenediamine dihydroiodide, cobalt carbonate, vitamin E supplement, thiamine mononitrate

WARNING: This product, which contains added copper, should not be fed to sheep or related species that have a low tolerance to copper.

This product has been formulated specifically for beef and is not intended for other species.

FEEDING DIRECTIONS:

Performance Elite Creep Feed Cattle Ration R40 is formulated for creep feeding calves from birth to weaning. Feed at the rate of 2.5 to 10.0 pounds per head per day to provide 50 to 200 milligrams Monensin per head per day. During the first 5 days of feeding, cattle should receive no more than 5.0 pounds (100 milligrams) Monensin per head per day. Over consumption may result in digestive disorders Remove spoiled feed regularly. Provide clean, fresh water at all times. Feed plain salt free-choice.

CAUTION: Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and the amount of refusals fed should be taken into consideration to prevent Monensin overdosing. A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

This feed does not contain urea, ruminant meat and bone meal, chicken litter, or restaurant grease.

Seller warrants only that this feed conforms to the description on the label and is suitable for the purposes set forth when used in accordance with the directions for use on said label.

Net weight 50 lb. (22.68 kg) or bulk invoice

MANUFACTURED BY:
PERFORMANCE FEEDS

2784 Bethany Road
Red Boiling Springs, TN 37150
(866) 699-5080
(615) 699-5080

LoadSheet #



PERFORMANCE ELITE CREEP FEED CATTLE RATION R40

MEDICATED

Calves-Birth to Weaning

Increased rate of weight gain in cattle (pastured slaughter, stocker, and feeder cattle, and dairy and beef replacement heifers).

ACTIVE DRUG INGREDIENT

MONENSIN.....40 g/ton

GUARANTEED ANALYSIS

CRUDE PROTEIN, min... 15.0 %	SALT, min.....0.1 %
CRUDE FAT, min..... 3.0 %	SALT, max..... 0.6 %
CRUDE FIBER, max..... 16.0 %	POTASSIUM, min.....0.7 %
CALCIUM, min.....1.1 %	VITAMIN A, min.....4,000 IU/lb
CALCIUM, max.....1.6 %	VITAMIN D, min.....400 IU/lb
PHOSPHORUS, min.....0.4 %	VITAMIN E, min.....5 IU/lb

INGREDIENTS

Processed grain by-products, roughage products, grain products, calcium carbonate, molasses products, salt, pork fat (preserved with BHA), zinc sulfate, manganese sulfate, ferrous sulfate, vitamin A supplement, vitamin D supplement, copper chloride, copper sulfate, sodium selenite, ethylenediamine dihydroiodide, cobalt carbonate, vitamin E supplement, thiamine mononitrate

WARNING: This product, which contains added copper, should not be fed to sheep or related species that have a low tolerance to copper.

This product has been formulated specifically for beef and is not intended for other species.

FEEDING DIRECTIONS:

Performance Elite Creep Feed Cattle Ration R40 is formulated for creep feeding calves from birth to weaning. Feed at the rate of 2.5 to 10.0 pounds per head per day to provide 50 to 200 milligrams Monensin per head per day. During the first 5 days of feeding, cattle should receive no more than 5.0 pounds (100 milligrams) Monensin per head per day. Over consumption may result in digestive disorders Remove spoiled feed regularly. Provide clean, fresh water at all times. Feed plain salt free-choice.

CAUTION: Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and the amount of refusals fed should be taken into consideration to prevent Monensin overdosing. A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

This feed does not contain urea, ruminant meat and bone meal, chicken litter, or restaurant grease.

Seller warrants only that this feed conforms to the description on the label and is suitable for the purposes set forth when used in accordance with the directions for use on said label.

Net weight 50 lb. (22.68 kg) or bulk invoice

MANUFACTURED BY:
PERFORMANCE FEEDS

2784 Bethany Road
Red Boiling Springs, TN 37150
(866) 699-5080
(615) 699-5080

LoadSheet #



PERFORMANCE ELITE CREEP FEED CATTLE RATION R40

MEDICATED

Calves-Birth to Weaning

Increased rate of weight gain in cattle (pastured slaughter, stocker, and feeder cattle, and dairy and beef replacement heifers).

ACTIVE DRUG INGREDIENT

MONENSIN.....40 g/ton

GUARANTEED ANALYSIS

CRUDE PROTEIN, min... 15.0 %	SALT, min.....0.1 %
CRUDE FAT, min..... 3.0 %	SALT, max..... 0.6 %
CRUDE FIBER, max..... 16.0 %	POTASSIUM, min.....0.7 %
CALCIUM, min.....1.1 %	VITAMIN A, min.....4,000 IU/lb
CALCIUM, max.....1.6 %	VITAMIN D, min.....400 IU/lb
PHOSPHORUS, min.....0.4 %	VITAMIN E, min.....5 IU/lb

INGREDIENTS

Processed grain by-products, roughage products, grain products, calcium carbonate, molasses products, salt, pork fat (preserved with BHA), zinc sulfate, manganese sulfate, ferrous sulfate, vitamin A supplement, vitamin D supplement, copper chloride, copper sulfate, sodium selenite, ethylenediamine dihydroiodide, cobalt carbonate, vitamin E supplement, thiamine mononitrate

WARNING: This product, which contains added copper, should not be fed to sheep or related species that have a low tolerance to copper.

This product has been formulated specifically for beef and is not intended for other species.

FEEDING DIRECTIONS:

Performance Elite Creep Feed Cattle Ration R40 is formulated for creep feeding calves from birth to weaning. Feed at the rate of 2.5 to 10.0 pounds per head per day to provide 50 to 200 milligrams Monensin per head per day. During the first 5 days of feeding, cattle should receive no more than 5.0 pounds (100 milligrams) Monensin per head per day. Over consumption may result in digestive disorders Remove spoiled feed regularly. Provide clean, fresh water at all times. Feed plain salt free-choice.

CAUTION: Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and the amount of refusals fed should be taken into consideration to prevent Monensin overdosing. A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

This feed does not contain urea, ruminant meat and bone meal, chicken litter, or restaurant grease.

Seller warrants only that this feed conforms to the description on the label and is suitable for the purposes set forth when used in accordance with the directions for use on said label.

Net weight 50 lb. (22.68 kg) or bulk invoice

MANUFACTURED BY:
PERFORMANCE FEEDS

2784 Bethany Road
Red Boiling Springs, TN 37150
(866) 699-5080
(615) 699-5080

LoadSheet #

