

Performance Safe-Guard® Dewormer

DAIRY AND BEEF CATTLE FEED FENBENDAZOLE TYPE C MEDICATED FEED

INDICATIONS

For the removal and control of:

Lungworms: (*Dictyocaulus viviparus*)

Stomach worms: Barberpole worms (*Haemonchus contortus*), brown stomach worms (*Ostertagia ostertagi*), small stomach worms (*Trichostrongylus axei*)

Intestinal worms: Hookworms (*Bunostomum phlebotomum*), thread-necked intestinal worms (*Nematodirus helvetianus*), small intestinal worms (*Cooperia punctata* & *C. oncophora*)

Bankrupt worms: (*Trichostrongylus colubriformis*)

Nodular worms: (*Oesophagostomum radiatum*)

ACTIVE DRUG INGREDIENTS

Fenbendazole443.4 g/ton

GUARANTEED ANALYSIS

CRUDE PROTEIN, min.....14.0 %
CRUDE FAT, min.....2.0 %
CRUDE FIBER, max.....25.0 %
ADF, max.....30.0 %
CALCIUM, min.....0.1 %
CALCIUM, max.....0.6 %
PHOSPHORUS, min.....0.4 %
POTASSIUM, min.....0.8 %

INGREDIENTS

Processed grain by-products, roughage products, forage products, mineral oil, and preserved with propionic acid.

FEEDING DIRECTIONS

Feed as the sole ration for one day. Feed at the rate of 1.02 pounds per 100 pounds of body weight to provide 2.27 milligrams fenbendazole per pound body weight.

CAUTION: All dewormers require accurate dosing for best results. Following the use of any dewormer, effectiveness of treatment should be monitored (for example, with the use of a fecal egg count reduction test or another appropriate method). A decrease of effectiveness over time as calculated by fecal egg count reduction tests may indicate the development of resistance to the dewormer administered. The parasite management plan should be adjusted accordingly based on regular monitoring.

WARNING:

RESIDUE WARNING: Cattle must not be slaughtered within 13 days following last treatment. For dairy cattle the milk discard time is zero hours. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

CONSULT YOUR VETERINARIAN FOR ASSISTANCE IN THE DIAGNOSIS, TREATMENT AND CONTROL OF PARASITISM

MANUFACTURED BY:

PERFORMANCE FEEDS

Load
Sheet#
212594
2784 Bethany Road
Red Boiling Springs, TN 37150
(866) 699-5080
(615) 699-5080

Net weight 50 lb. (22.68 kg) or bulk invoice

NADA # 131-675, Approved by FDA

Safe-Guard® is a registered trademark of Intervet Inc., d/b/a Merck Animal Health, a subsidiary of Merck & Co. Inc., All rights reserved.



Lot No.

Performance Safe-Guard® Dewormer

DAIRY AND BEEF CATTLE FEED FENBENDAZOLE TYPE C MEDICATED FEED

INDICATIONS

For the removal and control of:

Lungworms: (*Dictyocaulus viviparus*)

Stomach worms: Barberpole worms (*Haemonchus contortus*), brown stomach worms (*Ostertagia ostertagi*), small stomach worms (*Trichostrongylus axei*)

Intestinal worms: Hookworms (*Bunostomum phlebotomum*), thread-necked intestinal worms (*Nematodirus helvetianus*), small intestinal worms (*Cooperia punctata* & *C. oncophora*)

Bankrupt worms: (*Trichostrongylus colubriformis*)

Nodular worms: (*Oesophagostomum radiatum*)

ACTIVE DRUG INGREDIENTS

Fenbendazole443.4 g/ton

GUARANTEED ANALYSIS

CRUDE PROTEIN, min.....14.0 %
CRUDE FAT, min.....2.0 %
CRUDE FIBER, max.....25.0 %
ADF, max.....30.0 %
CALCIUM, min.....0.1 %
CALCIUM, max.....0.6 %
PHOSPHORUS, min.....0.4 %
POTASSIUM, min.....0.8 %

INGREDIENTS

Processed grain by-products, roughage products, forage products, mineral oil, and preserved with propionic acid.

FEEDING DIRECTIONS

Feed as the sole ration for one day. Feed at the rate of 1.02 pounds per 100 pounds of body weight to provide 2.27 milligrams fenbendazole per pound body weight.

CAUTION: All dewormers require accurate dosing for best results. Following the use of any dewormer, effectiveness of treatment should be monitored (for example, with the use of a fecal egg count reduction test or another appropriate method). A decrease of effectiveness over time as calculated by fecal egg count reduction tests may indicate the development of resistance to the dewormer administered. The parasite management plan should be adjusted accordingly based on regular monitoring.

WARNING:

RESIDUE WARNING: Cattle must not be slaughtered within 13 days following last treatment. For dairy cattle the milk discard time is zero hours. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

CONSULT YOUR VETERINARIAN FOR ASSISTANCE IN THE DIAGNOSIS, TREATMENT AND CONTROL OF PARASITISM

MANUFACTURED BY:

PERFORMANCE FEEDS

Load
Sheet#
212594
2784 Bethany Road
Red Boiling Springs, TN 37150
(866) 699-5080
(615) 699-5080

Net weight 50 lb. (22.68 kg) or bulk invoice

NADA # 131-675, Approved by FDA

Safe-Guard® is a registered trademark of Intervet Inc., d/b/a Merck Animal Health, a subsidiary of Merck & Co. Inc., All rights reserved.



Lot No.

Performance Safe-Guard® Dewormer

DAIRY AND BEEF CATTLE FEED FENBENDAZOLE TYPE C MEDICATED FEED

INDICATIONS

For the removal and control of:

Lungworms: (*Dictyocaulus viviparus*)

Stomach worms: Barberpole worms (*Haemonchus contortus*), brown stomach worms (*Ostertagia ostertagi*), small stomach worms (*Trichostrongylus axei*)

Intestinal worms: Hookworms (*Bunostomum phlebotomum*), thread-necked intestinal worms (*Nematodirus helvetianus*), small intestinal worms (*Cooperia punctata* & *C. oncophora*)

Bankrupt worms: (*Trichostrongylus colubriformis*)

Nodular worms: (*Oesophagostomum radiatum*)

ACTIVE DRUG INGREDIENTS

Fenbendazole443.4 g/ton

GUARANTEED ANALYSIS

CRUDE PROTEIN, min.....14.0 %
CRUDE FAT, min.....2.0 %
CRUDE FIBER, max.....25.0 %
ADF, max.....30.0 %
CALCIUM, min.....0.1 %
CALCIUM, max.....0.6 %
PHOSPHORUS, min.....0.4 %
POTASSIUM, min.....0.8 %

INGREDIENTS

Processed grain by-products, roughage products, forage products, mineral oil, and preserved with propionic acid.

FEEDING DIRECTIONS

Feed as the sole ration for one day. Feed at the rate of 1.02 pounds per 100 pounds of body weight to provide 2.27 milligrams fenbendazole per pound body weight.

CAUTION: All dewormers require accurate dosing for best results. Following the use of any dewormer, effectiveness of treatment should be monitored (for example, with the use of a fecal egg count reduction test or another appropriate method). A decrease of effectiveness over time as calculated by fecal egg count reduction tests may indicate the development of resistance to the dewormer administered. The parasite management plan should be adjusted accordingly based on regular monitoring.

WARNING:

RESIDUE WARNING: Cattle must not be slaughtered within 13 days following last treatment. For dairy cattle the milk discard time is zero hours. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

CONSULT YOUR VETERINARIAN FOR ASSISTANCE IN THE DIAGNOSIS, TREATMENT AND CONTROL OF PARASITISM

MANUFACTURED BY:

Load
Sheet#
212594
2784 Bethany Road
Red Boiling Springs, TN 37150
(866) 699-5080
(615) 699-5080

Net weight 50 lb. (22.68 kg) or bulk invoice

NADA # 131-675, Approved by FDA

Safe-Guard® is a registered trademark of Intervet Inc., d/b/a Merck Animal Health, a subsidiary of Merck & Co. Inc., All rights reserved.



Lot No.